

FREE CONSTRUCTION OF CAT(1) SPACES

KOICHI NAGANO

ABSTRACT. We construct several non-trivial examples of CAT(1) spaces by using the idea of free construction.

1. INTRODUCTION

A spherical (resp. Euclidean) building G of dimension n has plenty of apartments; more precisely, for any two points in G there exists an isometric embedding from the standard unit n -sphere (resp. the Euclidean n -space) to G whose image contains the given two points. Spherical or Euclidean buildings satisfy other rigid metric properties in the context of the geometry of CAT(1) spaces (cf. [KL]). In [CL, L, BL], we see several metric characterizations of buildings.

We say that a CAT(1) space is n -round if any two points are contained in an isometrically embedded unit n -sphere. For instance, every spherical building of dimension n is n -round. Every $\pi_{\mathbb{R}}$ -gon in the sense of [B] is 1-round. Every hemispherex of dimension $n \geq 2$ in [BB] is $(n - 1)$ -round (Example 2.4). In general, every n -round space is geodesically complete and $(n - 1)$ -round, and it has diameter π .

A geodesically complete $\mathbb{R}$ -tree with no graph structure has plenty of lines. This example suggests that only from the property of plenty of apartments one can not derive the rigid structure of buildings. Nevertheless, it seems to be non-trivial to see whether an n -round space of dimension n without building structure exists. The main purpose of this paper is to construct an n -round space of dimension n that is not a spherical building.

The Reshetnyak gluing theorem ([R], cf. Chapter II.11 in [BH]) helps us to obtain various examples of CAT(1) spaces. For our purpose, we need another idea. The purpose is achieved by mixing the idea of gluing with that of free construction. The idea of free construction was used by Tits [T] to obtain generalized polygons with large automorphism group.

Date: August 27, 2005.

2000 Mathematics Subject Classification. 53C20.

Key words and phrases. CAT(1) space.

Partially supported by the 2004 JSPS Postdoctoral Fellowships for Research Abroad.

For a CAT(1) space X , we denote by $\dim X$ the geometric dimension of X in the sense of [K]. By using the idea of free construction, we obtain the following:

Theorem 1.1. *Let $(X_\lambda)_{\lambda \in \Lambda}$ be a family of CAT(1) spaces of diameter $\leq \pi$. Then for any n there exists an n -round space Y such that*

- (1) $\dim Y = \max\{n, \sup_{\lambda \in \Lambda} \dim X_\lambda\}$;
- (2) *all points in Y are non-manifold points;*
- (3) *each X_λ is isometrically embedded in Y .*

In particular, there exists an n -round space of dimension n that is not a spherical building.

By our free construction, we also see the following. For a fixed $l \geq \pi$, let $(X_\lambda)_{\lambda \in \Lambda}$ be a family of CAT(1) spaces of diameter $\leq l$. Then for any n there exists a CAT(1) space in which any two points are contained in an isometrically embedded standard n -sphere of diameter l , and in which each X_λ is isometrically embedded (Theorem 3.5).

Next we consider the following condition. We say that two points in a metric space are *antipodes* if they have distance $\geq \pi$. A CAT(1) space is *n -rolling* if it has at least one pair of antipodes and if any antipodes are contained in an isometrically embedded unit n -sphere. Every n -round space is n -rolling. The unit n -hemisphere is $(n-1)$ -rolling. Every n -rolling space is geodesic and $(n-1)$ -rolling, and it has diameter π . If an n -rolling space has dimension n , then it is geodesically complete (cf. [BL]). It was proved in [BL] that if an n -rolling space of dimension n contains a relatively compact open set, then it is a spherical building.

A CAT(1) space has dimension 1 if and only if it is locally an $\mathbb{R}$ -tree of injectivity radius $\geq \pi$. A CAT(1) space of dimension 1 is 1-rolling if and only if it is 1-round. In general, we have:

Theorem 1.2. *For any $n \geq 2$, there exists an n -rolling space of dimension n that is not n -round.*

In fact, for any $n \geq 2$, such n -rolling spaces of dimension n can be obtained abundantly (Proposition 3.4).

We remark that the free construction discussed below also works in the CAT(κ) setting. The details are omitted in this paper.

We mention the relation between our notion and the asymptotic geometry of CAT(0) spaces. If H is a locally compact, geodesically complete CAT(0) space of dimension n , and if its Tits ideal boundary is $(n-1)$ -rolling, then H is a Euclidean building. This is substantially a corollary of the works of [CL, L, BL] (cf. Section 7 in [BL]).

2. PRELIMINARIES

2.1. CAT(1) spaces. A minimizing geodesic in a metric space means a length-minimizing curve joining two points. Any single point is also assumed to be a minimizing geodesic. A metric space is said to be

geodesic if any two points can be joined by a minimizing geodesic. A locally geodesic space is *geodesically complete* if any minimizing geodesic is contained in a bi-infinite locally minimizing geodesic.

We denote by $\mathbb{S}^n$ the standard unit n -sphere.

A complete metric space X with distance d is said to be CAT(1) if any two points $x, y \in X$ with $d(x, y) < \pi$ can be joined by a minimizing geodesic and if any geodesic triangle in X with perimeter $< 2\pi$ is not thicker than its comparison triangle in $\mathbb{S}^2$. If a CAT(1) space X is geodesically complete, then for any $x \in X$ we have a point $y \in X$ with $d(x, y) \geq \pi$. We refer to [BH] for the basics of CAT(1) spaces and the historical remarks in geometry of spaces with curvature bounded above in the sense of Alexandrov.

We first quote the Reshetnyak gluing theorem in the following form:

Theorem 2.1. ([R], cf. Chapter II.11 in [BH]) *Let $(X_\lambda)_{\lambda \in \Lambda}$ be a family of CAT(1) spaces with closed subspaces $C_\lambda \subset X_\lambda$, and X a CAT(1) space with a closed subspace $C \subset X$. Assume that*

- (1) *we have an isometry between C and C_λ ;*
- (2) *C is a CAT(1) space.*

Let $Y := X \sqcup (\bigsqcup_{\lambda \in \Lambda} X_\lambda) / \sim$ be the quotient metric space obtained by gluing X_λ to X along $C = C_\lambda$ for each λ . Then Y is a CAT(1) space.

Remark that a similar statement holds in the CAT(κ) setting. We use this gluing theorem only for the cases where C is a minimizing geodesic of length $\leq \pi$ (Subsection 3.2) and where C are antipodes (Subsection 3.3).

Let X be a CAT(1) space. For $x, y, z \in X$ with $d(x, y), d(x, z) < \pi$, the angle at x between y and z is denoted by $\angle_x(y, z)$. For $x \in X$, the space of directions at x equipped with the interior distance induced from $\angle_x$ is denoted by $\Sigma_x X$. This is also a CAT(1) space ([N]). The *geometric dimension* $\dim X$ of X is defined inductively as follows ([K]): If X is discrete, then $\dim X := 0$; in the other case,

$$\dim X := 1 + \sup_{x \in X} \dim \Sigma_x X.$$

Some geometric study of the geometric dimension can be seen in [K].

We say that a point $x \in X$ is *strongly singular in X* if $\Sigma_x X$ has at least two connected components. A manifold point means a point that has a neighborhood homeomorphic to a Euclidean space. We denote by $S(X)$ the set of all non-manifold points in X . Every manifold point in a CAT(1) space has the space of directions consisting of a single connected component. If x is strongly singular in X , then $x \in S(X)$.

2.2. Poles. Let X be a geodesically complete CAT(1) space of dimension ≥ 2 . A point $p \in X$ with $\dim \Sigma_p X \geq 1$ is called a *pole* if the diameter of $\Sigma_p X$ is greater than π . Remark that if X is n -round for $n \geq 2$, then it has no pole.

Example 2.2. For $n \geq 2$, the spherical suspension over a standard $(n-1)$ -sphere of diameter $> \pi$ is a geodesically complete CAT(1) space of dimension n and diameter π . The suspension points are poles.

Example 2.3. Let $n \geq 2$, and let $p, q \in \mathbb{S}^n$ be antipodes. Take some antipodes $\bar{p}, \bar{q}$ in another unit n -sphere $\mathbb{S}^n$. Let X be the quotient metric space $\mathbb{S}^n \sqcup \mathbb{S}^n / \sim$ made by attaching $\mathbb{S}^n$ to $\mathbb{S}^n$ at $p = \bar{p}$ and $q = \bar{q}$. This is a geodesically complete CAT(1) space of dimension n and diameter π . The antipodes p and q in X are poles.

Example 2.4. Let $n \geq 2$. A spherical simplicial complex is called an n -hemispherex if it is obtained from $\mathbb{S}^n$ by attaching unit n -hemispheres along great-hyperspheres such that no pair of antipodes in the original $\mathbb{S}^n$ belongs to all hyperspheres ([BB]). Every n -hemispherex is $(n-1)$ -round, not n -rolling, and has a pole.

2.3. Ultralimits. We review the notion of ultralimits, which goes back to [DW] (cf. Chapter I.5 in [BH]).

We fix a non-principal ultrafilter ω on $\mathbb{N}$. Let $(Y_i)_{i \in \mathbb{N}}$ be a sequence of metric spaces with distances d_i and basepoints p_i . We denote by Y_ω^0 the set of all sequences (y_i) such that $y_i \in Y_i$ and $d_i(p_i, y_i)$ is uniformly bounded. Let $d_\omega: Y_\omega^0 \times Y_\omega^0 \rightarrow [0, \infty)$ be the function defined as $d_\omega((y_i), (z_i)) := \omega\text{-}\lim d_i(y_i, z_i)$, where $\omega\text{-}\lim d_i$ is the ultralimit of the maps $d_i: Y_i \times Y_i \rightarrow [0, \infty)$. We define $\omega\text{-}\lim Y_i$ as the quotient metric space $(Y_\omega^0, d_\omega) / d_\omega = 0$, called the *ultralimit of (Y_i) with respect to ω* . Note that if the diameters of Y_i are uniformly bounded, then $\omega\text{-}\lim Y_i$ does not depend on the choices of the basepoints p_i . If each Y_i is complete (resp. geodesically complete), then $\omega\text{-}\lim Y_i$ is again complete (resp. geodesically complete). If we have a metric space X and isometric embeddings $f_i: X \rightarrow Y_i$, then we have a natural isometric embedding $f_\omega: X \rightarrow \omega\text{-}\lim Y_i$ defined by $f_\omega(x) := (f_i(x))$. If each Y_i is CAT(1), then $\omega\text{-}\lim Y_i$ is again CAT(1); moreover, if $\dim Y_i \leq n$ for each i , then $\dim(\omega\text{-}\lim Y_i) \leq n$ (cf. [L]).

Let Y be a metric space, and ω a non-principal ultrafilter on $\mathbb{N}$. We call the ultralimit $\omega\text{-}\lim Y$ for the constant sequence (Y) the *ultra-completion of Y with respect to ω* . By definition, Y can be naturally isometrically embedded in $\omega\text{-}\lim Y$. In the sequel, we use only ultra-completions of metric spaces.

3. CONSTRUCTION

We prove Theorems 1.1 and 1.2.

3.1. Truncated inductive limits. For metric spaces X and Y , we write $X < Y$ if we have an isometric embedding from X to Y . We say that a map $f: X \rightarrow Y$ is a π -truncated isometric embedding if for any $x, y \in X$ with $d(x, y) \leq \pi$ we have $d(f(x), f(y)) = d(x, y)$. We write $X \prec Y$ if we have a π -truncated isometric embedding from X to Y .

Let $Y_0 \prec Y_1 \prec \cdots \prec Y_{i-1} \prec \cdots$ be an inductive system of metric spaces Y_{i-1} with distances d_{i-1} , $i \in \mathbb{N}$, and π -truncated isometric embeddings from Y_{i-1} to Y_i . We denote by $Y_\infty := \bigcup_{i \in \mathbb{N}} Y_{i-1}$ the inductive limit space with the distance d defined by $d|_{Y_{i-1} \times Y_{i-1}} := \min\{\pi, d_{i-1}\}$. We call Y_∞ the π -truncated inductive limit. If Y_i is geodesically complete for each i , then so is Y_∞ . We need the following later on.

Lemma 3.1. *For a sequence of CAT(1) spaces Y_{i-1} of $\dim Y_{i-1} \leq m$ with $Y_{i-1} \prec Y_i$, $i \in \mathbb{N}$, let $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$ be the π -truncated inductive limit, and $\omega\text{-lim } Y_\infty$ the ultracompletion of Y_∞ . Then $\omega\text{-lim } Y_\infty$ is a CAT(1) space of $\dim(\omega\text{-lim } Y_\infty) \leq m$. Moreover, if in addition Y_i is geodesically complete for each i , then we have the following:*

- (1) *if each point in Y_∞ is strongly singular in some Y_i , then each point in $\omega\text{-lim } Y_\infty$ is again strongly singular;*
- (2) *if any two points in Y_∞ are contained in an isometrically embedded unit n -sphere, then $\omega\text{-lim } Y_\infty$ is n -round;*
- (3) *if any antipodes in Y_∞ are contained in an isometrically embedded unit n -sphere, then $\omega\text{-lim } Y_\infty$ is n -rolling.*

Proof. Since each Y_i is CAT(1), $\omega\text{-lim } Y_\infty$ is again CAT(1). The dimension does not increase under the ultralimits procedures (cf. [L]). Hence we have $\dim(\omega\text{-lim } Y_\infty) \leq m$. Observing the ultralimits of isometrically embedded unit n -spheres in $\omega\text{-lim } Y_\infty$, we have (2) and (3). We show (1). Suppose the contrary, i.e., we have a point (y) in $\omega\text{-lim } Y_\infty$ such that $\Sigma_{(y)}(\omega\text{-lim } Y_\infty)$ has a unique connected component, where $y \in Y_\infty$. Then y is strongly singular in some Y_i . Hence $\Sigma_y Y_i$ has at least two components. Since each Y_i is geodesically complete, $\Sigma_{(y)}(\omega\text{-lim } Y_\infty)$ must have at least two components. This is a contradiction. $\square$

3.2. Round spaces. We prove Theorem 1.1. For a CAT(1) space X , we denote by Γ_X the set of all minimizing geodesics in X of length $\leq \pi$.

By using the idea of free construction, we first show the following:

Proposition 3.2. *Let X be a CAT(1) space of diameter $\leq \pi$. Then for any n there exists an n -round space Y with $\dim Y = \max\{n, \dim X\}$, $Y = S(Y)$, and $X < Y$.*

Proof. For a fixed n , let $m := \max\{n, \dim X\}$. We construct a CAT(1) space Y inductively as follows. Put $Y_0 := X$. Let Y_{i-1} be a CAT(1) space of $\dim Y_{i-1} \leq m$. Set $\Gamma_{i-1} := \Gamma_{X_{i-1}}$. For each $\gamma \in \Gamma_{i-1}$, we take a minimizing geodesic $\bar{\gamma}$ in $\mathbb{S}^n = \mathbb{S}^n(\gamma)$ of the same length. We define Y_i as the quotient metric space $Y_{i-1} \sqcup (\bigsqcup_{\gamma \in \Gamma_{i-1}} \mathbb{S}^n(\gamma)) / \sim$ made by gluing $\mathbb{S}^n(\gamma)$ to Y_{i-1} along $\gamma = \bar{\gamma}$. For each $i \in \mathbb{N}$, Y_i is a geodesically complete, geodesic space. By the Reshetnyak gluing theorem, Y_i is CAT(1). It follows that $\dim Y_i = m$. Let $f_i: Y_{i-1} \rightarrow Y_i$ be the inclusion. This is a π -truncated isometric embedding, and hence $Y_{i-1} \prec Y_i$. For any $x, y \in Y_{i-1}$, $f_i(x)$ and $f_i(y)$ are contained in an embedded unit n -sphere.

For any $z \in Y_{i-1}$, $f_i(z)$ is strongly singular in Y_i . Let $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$ be the π -truncated inductive limit. Each point in Y_∞ is strongly singular in some Y_i , and any two points are contained in an embedded unit n -sphere. For a given non-principal ultrafilter ω on $\mathbb{N}$, let $Y := \omega\text{-lim } Y_\infty$ be the ultracompletion of Y_∞ . By Lemma 3.1, Y is an n -round space with $\dim Y = m$, $Y = S(Y)$, and $X < Y$. $\square$

We notice that each Y_i has diameter $> \pi$. The space Y_∞ does not depend on the choices of minimizing geodesics in $\mathbb{S}^n$.

Remark 3.3. We recall that CAT(1) spaces in this paper are assumed to be complete. This seems to be natural since almost all significant CAT(1) spaces are complete.

In the proof of Proposition 3.2, if one does not require the completeness for the definition of CAT(1) spaces, then Y_∞ is a space with the desired property. To obtain the completeness, we need to take the ultracompletion $\omega\text{-lim } Y_\infty$. The usual completion $\overline{Y}_\infty$ of Y_∞ does not work for our purpose since we can not see whether embedding unit n -spheres in Y_∞ converge to the desired spheres in $\overline{Y}_\infty$.

Now we prove Theorem 1.1.

Proof. Let $(X_\lambda)_{\lambda \in \Lambda}$ be a family of CAT(1) spaces of diameter $\leq \pi$. We fix n . For each X_λ , by Proposition 3.2 we have an n -round space X'_λ with $\dim X'_\lambda = \max\{n, \dim X_\lambda\}$, $X'_\lambda = S(X'_\lambda)$, and $X_\lambda < X'_\lambda$.

Let $m := \max\{n, \sup_{\lambda \in \Lambda} \dim X_\lambda\}$. We construct a space Y inductively in the following way. Put $Y_0 := X'_{\lambda_0}$ for some λ_0 . Let Y_{i-1} be a geodesically complete, geodesic CAT(1) space of $\dim Y_{i-1} \leq m$. Set $\Gamma_{i-1} := \Gamma_{Y_{i-1}}$. For each $(\gamma, \lambda) \in \Gamma_{i-1} \times \Lambda$, take a minimizing geodesic $\overline{\gamma}$ in $X'_\lambda = X'_\lambda(\gamma)$ of the same length. Let

$$Y_i := Y_{i-1} \sqcup \left(\bigsqcup_{(\gamma, \lambda) \in \Gamma_{i-1} \times \Lambda} X'_\lambda(\gamma) \right) / \sim$$

be the quotient metric space made by gluing $X'_\lambda(\gamma)$ to Y_{i-1} along $\gamma = \overline{\gamma}$ for each λ . The space Y_i is a geodesically complete, geodesic CAT(1) space of $\dim Y_i = m$. Then $Y_{i-1} \prec Y_i$ by the inclusion $f_i: Y_{i-1} \rightarrow Y_i$, and $X'_\lambda < Y_i$ for each λ . For any $x, y \in Y_{i-1}$, $f_i(x)$ and $f_i(y)$ are contained in an embedded unit n -sphere. For any $z \in Y_{i-1}$, $f_i(z)$ is strongly singular in Y_i . For a given non-principal ultrafilter ω on $\mathbb{N}$, let $Y := \omega\text{-lim } Y_\infty$ be the ultracompletion of the π -truncated inductive limit $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_{i-1}$. By Lemma 3.1, Y is an n -round space with $\dim Y = m$ and $Y = S(Y)$. Moreover, $X_\lambda < Y$ for each λ . Thus we obtain Theorem 1.1. $\square$

3.3. Rolling spaces. For a CAT(1) space X , we denote by A_X the subset of $X \times X$ consisting of all pairs of antipodes in X .

To prove Theorem 1.2, it suffices to show:

Proposition 3.4. *Let $n \geq 2$, and let X be a geodesically complete, geodesic CAT(1) space of dimension n and diameter π . Assume that X has a pole. Then there exists an n -rolling space Y of dimension n such that $Y = S(Y)$, $X < Y$, and Y is not n -round.*

Proof. We first construct a space Y as follows. Put $Y_0 := X$. Let Y_{i-1} be a geodesically complete, geodesic CAT(1) space of dimension n , and set $A_{i-1} := A_{Y_{i-1}}$. For each $(p, q) \in A_{i-1}$, take some antipodes $\bar{p}, \bar{q}$ in $\mathbb{S}^n = \mathbb{S}^n(p, q)$. Define Y_i as $Y_{i-1} \sqcup (\bigsqcup_{(p,q) \in A_{i-1}} \mathbb{S}^n(p, q) / \sim$ made by attaching $\mathbb{S}^n(p, q)$ to Y_{i-1} at $p = \bar{p}$ and $q = \bar{q}$. The space Y_i is a geodesically complete, geodesic CAT(1) space of dimension n . Then $Y_{i-1} \prec Y_i$ by the inclusion $f_i: Y_{i-1} \rightarrow Y_i$. For each $(x, y) \in A_{i-1}$, the points $f_i(x)$ and $f_i(y)$ are contained in an embedded unit n -sphere. For each i and for each $z \in Y_i$, $f_{i+1}(z)$ is strongly singular in Y_{i+1} . Let $Y_\infty = \bigcup_{i \in \mathbb{N}} Y_i$ be the π -truncated inductive limit. Each point in Y_∞ is strongly singular in some Y_i , and any antipodes in Y_∞ are contained in an embedded unit n -sphere. For a given non-principal ultrafilter ω on $\mathbb{N}$, let $Y := \omega\text{-lim } Y_\infty$ be the ultracompletion of Y_∞ . By Lemma 3.1, Y is an n -rolling space of dimension n with $Y = S(Y)$ and $X < Y$. Let $f: X \rightarrow Y$ be an isometric embedding, and $p \in X$ a pole. Then $f(p) \in Y$ is again a pole. Hence Y is not n -round since every n -round space has no pole. This completes the proof. $\square$

The CAT(1) spaces in Examples 2.2, 2.3, and 2.4 satisfy the assumption in Proposition 3.4. Hence we have proved Theorem 1.2. $\square$

3.4. Modification. For $l \geq \pi$, we say that a CAT(1) space is (n, l) -round if any two points are contained in an isometrically embedded standard n -sphere $\mathbb{S}_l^n$ of diameter l .

Let X be a CAT(1) space. For $l \geq \pi$, we denote by Γ_X^l the set of all minimizing geodesics in X of length $\leq l$. In the proof of Theorem 1.1, by replacing $\mathbb{S}^n$ (resp. Γ_X) with $\mathbb{S}_l^n$ (resp. Γ_X^l), and by considering l -truncated isometric embeddings and l -truncated inductive limits instead of π -truncated ones, we obtain:

Theorem 3.5. *For a fixed $l \geq \pi$, let $(X_\lambda)_{\lambda \in \Lambda}$ be a family of CAT(1) spaces of diameter $\leq l$. Then for any n there exists an (n, l) -round space Y such that $\dim Y = \max\{n, \sup_{\lambda \in \Lambda} \dim X_\lambda\}$, $Y = S(Y)$, and $X_\lambda < Y$ for each $\lambda \in \Lambda$.*

Acknowledgments. I am grateful to Alexander Lytchak for his valuable comment, to Werner Ballmann for his comment on this work, and to Linus Kramer for his information on [B]. This paper was written during my stay at the University of Bonn. I would like to thank all geometers in Bonn for their great hospitality.

REFERENCES

- [BB] W. Ballmann and M. Brin, *Diameter rigidity of spherical polyhedra*, Duke Math. J. **122** (1999), 597–609.
- [BL] A. Balser and A. Lytchak, *Building-like spaces*, preprint, 2004; Math. ArXiv., math.MG/0410437.
- [B] C. D. Bennet, *Twin trees and λ_Λ -gons*, Trans. Amer. Math. Soc. **349** (1997), Number 5, 2069–2084.
- [BH] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Math. Wiss., Volume 319, Springer-Verlag, 1999.
- [CL] R. Charney and A. Lytchak, *Metric characterizations of spherical and Euclidean buildings*, Geom. Topol. **5** (2001), 521–550.
- [DW] L. van den Dries and A. J. Wilkie, *On Gromov’s theorem concerning groups of polynomial growth and elementary logic*, J. Algebra **89** (1984), 349–374.
- [K] B. Kleiner, *The local structure of length spaces with curvature bounded above*, Math. Z. **231** (1999), 409–456.
- [KL] B. Kleiner and B. Leeb, *Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings*, Inst. Hautes Études Sci. Publ. Math. **86** (1997), 115–197 (1998).
- [L] A. Lytchak, *Rigidity of spherical buildings and spherical joins*, preprint, to appear in Geom. Funct. Anal.
- [N] I. G. Nikolaev, *The tangent cone of an Alexandrov space of curvature $\leq K$* , manuscripta math. **86** (1995), 683–689.
- [R] Yu. G. Reshetnyak, *On the theory of spaces of curvature not greater than K* , Mat. Sb. **52** (1960), 789–798.
- [T] J. Tits, *Endliche Spiegelungsgruppen, die als Weylgruppen auftreten*, Invent. Math. **43**, (1977), 283–295.

E-mail address: nagano@math.tohoku.ac.jp / nagano@math.uni-bonn.de

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY
AOBA, SENDAI, MIYAGI, 980-8578, JAPAN

Current address: Mathematisches Institut, Universität Bonn
Berlingstraße 1, D-53115 Bonn, Germany